

# INSECTICIDE RESISTANCE IN INSECT VECTORS OF HUMAN DISEASE

---

Janet Hemingway and Hilary Ranson

*School of Biosciences, University of Wales Cardiff, P.O. Box 915, Cardiff, Wales CF1  
3TL; e-mail: sabjh@cardiff.ac.uk*

**Key Words** insecticide, mosquito, esterases, monooxygenases, glutathione S-transferases

■ **Abstract** Insecticide resistance is an increasing problem in many insect vectors of disease. Our knowledge of the basic mechanisms underlying resistance to commonly used insecticides is well established. Molecular techniques have recently allowed us to start and dissect most of these mechanisms at the DNA level. The next major challenge will be to use this molecular understanding of resistance to develop novel strategies with which we can truly manage resistance. State-of-the-art information on resistance in insect vectors of disease is reviewed in this context.

## INTRODUCTION

Insecticides play a central role in controlling major vectors of diseases such as mosquitoes, sandflies, fleas, lice, tsetse flies, and triatomid bugs. In 1955 the World Health Organization (WHO) assembly proposed the global eradication of the most prevalent vector-borne human disease, malaria, by the use of residual house-spraying of DDT. However, the insecticide euphoria soon ended and in 1976 WHO officially reverted from malaria eradication to malaria control. This marked shift from malaria eradication to primary health care was an emotive issue, eliciting a rapid and complete change of rhetoric from WHO (12). Several issues had prompted this switch, but a major cause of the change in policy was the appearance of DDT resistance in a broad range of the mosquito vectors. In 1975 WHO reported that 256 million people were living in areas where DDT and/or BHC resistance was undermining malaria control efforts. (This did not include the African region, where 90% of malaria occurs and where DDT resistance had already been noted in *Anopheles gambiae*, the major malaria vector.)

The resistance problems continued with the switch to newer insecticides such as the organophosphates, carbamates and pyrethroids. Operationally, many control programs have switched from blanket spraying of house interiors to focal use of insecticides on bednets. Focal spraying limits the insecticides of choice largely to pyrethroids due to the speed of kill required to protect the occupant of the

bednet and the safety margin needed for insecticides used in such close contact with people. Today the major emphasis in resistance research is on the molecular mechanisms of resistance and rational resistance management, with a view to controlling the development and spread of resistant vector populations. In Africa, WHO and the World Bank have instigated major new initiatives with other major donors and the scientific community internationally to “roll back” malaria. One major problem these initiatives are tackling is the presence of two developing foci of pyrethroid resistance in the most important African malaria vector, *An. gambiae*.

## SCALE OF THE PROBLEM

The amount of resistance in insect vector populations is dependent both on the volume and frequency of applications of insecticides used against them and the inherent characteristics of the insect species involved. Tsetse flies, for example, were controlled by wide-scale spraying of DDT for many years, but DDT resistance has never developed in this species. Another example of an insect vector exhibiting little or no resistance to insecticides is the triatomid bug. In both cases the major factor influencing insecticide resistance development is the life cycle of the insect pest, in particular the long life cycles for the bugs, and the production of very small numbers of young by the tsetse flies. In contrast, mosquitoes have all the characteristics suited to rapid resistance development, including short life cycles with abundant progeny.

## Mosquito Resistance

The major mosquito vectors span the *Culex*, *Aedes*, and *Anopheles* genera. *Culex* are the major vectors of filariasis and Japanese encephalitis, *Aedes* of dengue and dengue hemorrhagic fever, and *Anopheles* of malaria. The range of many of these species is not static. For example, several *Aedes* species recently extended their range in Asia and Latin America, leading to an increased risk of dengue in these areas.

DDT was first introduced for mosquito control in 1946. In 1947 the first cases of DDT resistance occurred in *Aedes tritaeniorhynchus* and *Ae. sollicitans* (15). Since then more than 100 mosquito species are reported as resistant to one or more insecticide, and more than 50 of these are anophelines (113). Insecticides used for malaria control have included -BHC, organophosphorus, carbamate, and pyrethroid insecticides, with the latter now taking increasing market share for both indoor residual spraying and large-scale insecticide-impregnated bednet programs. Other insecticide groups, such as the benzylphenyl ureas and Bti, have had limited use against mosquitoes. Resistance has tended to follow the switches of insecticides. Resistance to -BHC/dieldrin is widespread despite the lack of use of these insecticides for many years. Organophosphate (OP) resistance, either in

the form of broad-spectrum OP resistance or malathion-specific resistance, occurs in the major vectors *An. culicifacies* (59), *An. stephensi* (30, 44), *An. albimanus* (3, 51), *An. arabiensis* (45) and *An. sacharovi* (54). *An. culicifacies* is recognized as a species complex (101): In Sri Lanka malathion resistance occurs in *An. culicifacies* species B, while in India resistance is in species B and C (59, 88). Species B in Sri Lanka is resistant to fenitrothion, which is independent of the malathion-specific resistance (58, 60), and is developing pyrethroid resistance (SHPP Karunaratne, personal communication). Organophosphorus insecticide resistance is widespread in all the major *Culex* vectors (53), and pyrethroid resistance occurs in *C. quinquefasciatus* (1, 7, 19). Pyrethroid resistance has been noted in *An. albimanus* (13), *An. stephensi* (107) and *An. gambiae* (18, 20, 112) among others, while carbamate resistance is present in *An. sacharovi* and *An. albimanus* (57). Pyrethroid resistance is widespread in *Ae. aegypti* (6, 48, 70) and cases of OP and carbamate resistance have also been recorded in this species (72, 76).

The development of pyrethroid resistance in *An. gambiae* is particularly important given the recent emphasis by the WHO and other organizations on the use of pyrethroid-impregnated bednets for malaria control. Two cases of pyrethroid resistance in *An. gambiae*, from the Ivory Coast and Kenya, are well documented (18, 112). The west African focus appears to be larger and has higher levels of resistance than that in east Africa.

### Sandfly Resistance

The peridomestic vectors of *Leishmania*, *Plebotomus papatasi*, *Lutzomyia longipalpis*, and *L. intermedia* are controlled primarily by insecticides throughout their range. The control of these sandflies is often a by-product of anti-malarial house-spraying. The only insecticide resistance reported to date in sandflies is to DDT in Indian *P. papatasi* (29).

### Head and Body Louse Resistance

The body louse *Pediculus humanus* has developed widespread resistance to organochlorines (16), is malathion resistant in parts of Africa (113), and has low-level resistance to pyrethroids in several regions (35). Resistance to organochlorine insecticides, such as DDT and lindane, has been recorded in the human head louse *Pediculus capitis* in Israel, Canada, Denmark, and Malaysia (16, 73, 113). Permethrin has been extensively used for head louse control since the early 1980s (77, 103). The first reports of control failure with this insecticide were in the early 1990s in Israel (77), the Czech Republic (97), and France (21).

### *Simulium* Resistance

Some cytospecies of the *Simulium damnosum* complex are vectors of onchocerciasis. In 1974 the Onchocerciasis Control Programme in west Africa established a long-term insecticide-based control program for this vector. Temephos resis-

tance occurred initially, prompting a switch to chlorphoxim, but resistance to this insecticide occurred within a year (49). Resistance in this species is currently being managed by a rotation of temephos, Bti, and permethrin, the insecticide usage being determined by the rate at which water is flowing in rivers forming the major breeding sites of these vectors.

## THE BIOCHEMISTRY OF RESISTANCE

### Insecticide Metabolism

Three major enzyme groups are responsible for metabolically based resistance to organochlorines, organophosphates, carbamates, and pyrethroids. DDT-dehydrochlorinase was first recognized as a glutathione S-transferase in the house fly, *Musca domestica* (23). It has been shown to have this role commonly in anopheline and *Aedes* mosquitoes (40, 85). Esterases are often involved in organophosphate, carbamate, and to a lesser extent, pyrethroid resistance. Monooxygenases are involved in the metabolism of pyrethroids, the activation and/or detoxication of organophosphorus insecticides and, to a lesser extent, carbamate resistance.

### Esterase-Based Resistance

The esterase-based resistance mechanisms have been studied most extensively at the biochemical and molecular level in *Culex* mosquitoes and the aphid *Myzus persicae*. Work is in progress on related and distinct esterase resistance mechanisms in a range of *Anopheles* and *Aedes* species. Broad-spectrum organophosphate resistance is conferred by the elevated esterases of *Culex*. All these esterases act by rapidly binding and slowly turning over the insecticide: They sequester rather than rapidly metabolize the pesticide (62).

Two common esterase loci, *est $\alpha$*  and *est $\beta$* , are involved alone or in combination in this type of resistance in *Culex* (109). In *C. quinquefasciatus* the most common elevated esterase phenotype involves two enzymes, *est $\alpha$ 2<sup>1</sup>* and *est $\beta$ 2<sup>1</sup>* (*A<sub>2</sub>* and *B<sub>2</sub>* on an earlier classification) (110). The classification of these esterases is based on their preferences for  $\alpha$ - or  $\beta$ -naphthyl acetate, their mobility on native polyacrylamide gels, and their nucleotide sequence (53). Smaller numbers of *C. quinquefasciatus* populations have elevated *est $\beta$ 1* alone, elevated *est $\alpha$ 1* alone or co-elevated *est $\beta$ 1* and *est $\alpha$ 3* (27, 53). When purified *est $\alpha$*  and *est $\beta$*  from the insecticide-susceptible PeISS strain were compared to various enzymes purified from resistant strains, up to 1000-fold differences among the inhibition-kinetic constants occurred for the oxon analogues of various OPs (63).

The superiority of insecticide binding in enzymes from the resistant strains suggests that there has been positive insecticide selection pressure to maintain elevation of favorable alleles of the esterases in insecticide-resistant insects. Although there are minor variations between the inhibition kinetics of the different elevated alleles, the reason why the *est $\alpha$ 2<sup>1</sup>/est $\beta$ 2<sup>1</sup>* phenotype is so common

(in more than 90% of resistant populations) compared to the other elevated esterase phenotypes is not obvious. This advantage may be linked to a third gene, which is co-elevated with esterases est $\alpha$ 2/est $\beta$ 2 but not with the other esterase phenotypes (50).

Metabolic studies on *Culex* homogenates suggests that increased rates of esterase-mediated metabolism plays little or no role in resistance. One exception to this is *C. tarsalis*, where two resistance mechanisms co-exist: one involving elevated sequestering esterases, the other involving non-elevated metabolically active esterases (120). In contrast to the situation in *Culex*, a number of *Anopheles* species have a non-elevated esterase mechanism that confers resistance specifically to malathion through increased rates of metabolism (44–46) (11, 68). In *An. stephensi* three esterases with malathion carboxylesterase activity have been isolated and characterized (47, 52).

### Glutathione S-Transferase-Based Resistance

Many studies have shown that insecticide-resistant insects have elevated levels of glutathione S-transferase activity in crude homogenates, which suggests a role for GSTs in resistance (37, 38). GSTs are dimeric multifunctional enzymes that play a role in detoxification of a large range of xenobiotics (86). The enzymes catalyze the nucleophilic attack of reduced glutathione (GSH) on the electrophilic centers of lipophilic compounds. Multiple forms of these enzymes have been reported for mosquitoes, house fly, *Drosophila*, sheep blow fly, and grass grub (22, 24, 105).

Two families of insect GST are recognized, and both appear to have a role in insecticide resistance in insects. In *Ae. aegypti* at least two GSTs are elevated in DDT-resistant insects (39, 41), while in *An. gambiae* a large number of different GSTs are elevated, some of which are class I GSTs (84, 85). The *Ae. aegypti* and *An. gambiae* GSTs in resistant insects are constitutively over-expressed. The GST-2 of *Ae. aegypti* is over-expressed in all tissues except the ovaries of resistant insects (39).

### Monoxygenase-Based Resistance

The monooxygenases are a complex family of enzymes found in most organisms, including insects. These enzymes are involved in the metabolism of xenobiotics and have a role in endogenous metabolism. The P450 monooxygenase are generally the rate-limiting enzyme step in the chain. These enzymes are important in adaptation of insects to toxic chemicals in their host plants. P450 monooxygenases are involved in the metabolism of virtually all insecticides, leading to activation of the molecule in the case of organophosphorus insecticides, or more generally to detoxification. P450 enzymes bind molecular oxygen and receive electrons from NADPH to introduce an oxygen molecule into the substrate.

Elevated monooxygenase activity is associated with pyrethroid resistance in *An. stephensi*, *An. subpictus*, *An. gambiae* (14, 55, 112), and *C. quinquefasciatus*

(65). Currently this enzyme system is poorly studied in insect vectors of disease. The nomenclature of the P450 superfamily is based on amino acid sequence homologies, with all families having the CYP prefix followed by a numeral for the family, a letter for the subfamily, and a numeral for the individual gene. To date insect P450s have been assigned to six families: five are insect-specific and one, CYP4, has sequence homologies with families in other organisms (8).

### Target-Site Resistance

The organophosphorus, carbamates, organochlorine, and pyrethroid insecticides all target the nervous system. Newer classes of insecticides are available for vector control, but the high cost of developing and registering new insecticides inevitably means that insecticides are developed initially for the agricultural market and then utilized for public health vector control, where their activities and safety profile are appropriate and where the market is sufficiently large to warrant the registration costs for public health use. Compounds targeting the nicotinic acetylcholine receptor have recently made this transition from agriculture into public health.

### Acetylcholinesterase

The organophosphates and carbamates target acetylcholinesterase (AChE). AChE hydrolyzes the excitatory neurotransmitter acetylcholine on the post-synaptic nerve membrane. Insect AChE has a substrate specificity intermediate between vertebrate AChE and butyrylcholinesterase. The predominant molecular form in insects is a globular amphiphilic dimer which is membrane-bound via a glycolipid anchor. Alterations in AChE in organophosphate- and carbamate-resistant insects result in a decreased sensitivity to inhibition of the enzyme by these insecticides (5, 51). The organophosphorus insecticides are converted to their oxon analogues via the action of monooxygenases before acting as AChE inhibitors. In *C. pipiens*, AChE1 and AChE2 differ in their substrate specificity, inhibitor sensitivity, and electrophoretic migration pattern (69). Only AChE1 appears to be involved in conferring insecticide resistance.

### GABA Receptors

Resistance to dieldrin was recorded in the 1950s, but the involvement of the GABA receptors in this resistance was not elucidated until the 1990s. The GABA receptor in insects is a heteromultimeric gated chloride-ion channel, a widespread inhibitory neurotransmission channel in the insect's central nervous system and in neuromuscular junctions (9). The insect GABA receptor is implicated as a site of action for pyrethroids and avermectins as well as cyclodienes. Studies showing that cyclodiene-resistant insects are resistant to picrotoxin and phenylpyrazole insecticides, and that the effect of ivermectin on cultured neurons can be reversed by picrotoxin pretreatment, suggest that these insecticides exert their effect by

interacting with the chloride ionophore associated with the insect GABA receptor (10, 62).

### Sodium Channels

The pharmacological effect of DDT and pyrethroids is to cause persistent activation of the sodium channels by delaying the normal voltage-dependent mechanism of inactivation (100). Insensitivity of the sodium channels to insecticide inhibition was first recorded in *Musca domestica* (32). In mosquitoes there have been many reports of suspected “kdr”-like resistance inferred from cross resistance between DDT and pyrethroids, which act on the same site within the sodium channel. These reports have been validated by electrophysiological measurements in *Ae. aegypti* and *An. stephensi* (48, 107).

## THE MOLECULAR BIOLOGY OF RESISTANCE

### Metabolic Mechanisms

Over-expression of enzymes capable of detoxifying insecticides or amino acid substitutions within these enzymes, which alter the affinity of the enzyme for the insecticide, can result in high levels of insecticide resistance. Increased expression of the genes encoding the major xenobiotic metabolizing enzymes are the most common cause of insecticide resistance in mosquitoes. These large enzyme families may contain multiple enzymes with broad overlapping substrate specificities, and there is a high probability that at least one member of the family will be capable of metabolizing one or more insecticides. Increased production of these enzymes may have a lower associated fitness cost than those associated with alterations in the structural genes because the primary function of the enzyme is not disrupted.

### Mutations in Structural Genes

In many cases of resistance caused by increased metabolism of the insecticide the exact genetic mechanism is not known. As yet no validated reports exist of mutations within detoxifying enzymes leading to resistance to insecticides in disease vectors. Two examples have been reported in non-vector species: because both of these mechanisms may be present in disease vectors, we describe them here.

Resistance to the organophosphate insecticide malathion is caused by a single amino acid substitution (Trp<sup>251</sup>-Leu) within the E3 esterase of the sheep blow fly, *Lucilia cuprina* (17). Malathion-resistant strains of *L. cuprina* have very low levels of activity with aliphatic esters that are conventionally used as stains for esterase activity (80). A similar phenotype has been observed in malathion-resistant strains of *An. stephensi*, *An. arabiensis*, and *An. culicifacies*. At least three enzymes are able to metabolize malathion in *An. stephensi* but it is not yet

known whether a point mutation similar to that described in *Lucilia* is responsible for malathion resistance in *Anopheles* (51).

A second distinct amino acid substitution (Gly<sup>137</sup>-Asp) within the active site of the *Lucilia* E3 isozyme confers broad cross resistance to many organophosphorus insecticides but not to malathion (79). This same mutation is present in OP-resistant strains of house fly.

## Gene Amplification

The metabolic resistance mechanism studied in most detail in insect disease vectors is the elevated esterase-based system in *Culex* (110, 111). In the three *Culex* species studied to date at the molecular level the homologous *estβ* gene is amplified in resistant insects (64, 111, 114, 115). Insecticide resistance via amplification of genes involved in their detoxification is common in several insects. The most common amplified esterase-based mechanism in *Culex* involves the co-amplification of two esterases, *estα2*<sup>1</sup> and *estβ2*<sup>1</sup> in *C. quinquefasciatus* and other members of the *C. pipiens* complex worldwide (109). Other strains of *C. quinquefasciatus* have *estα3* and *estβ1* co-amplified (27), while the TEM-R strain has amplified *estβ1* alone (75). Similarly, *C. tritaeniorhynchus* has a single amplified *estβ*, *Ctrestβ1* (64).

The *estα* and *estβ* genes have arisen as the result of a gene duplication event, which must have occurred prior to speciation within the *Culex* genus. The genes occur in a head-to-head arrangement approximately 1.7 Kb apart in susceptible insects (109). In resistant insects with *estα2*<sup>1</sup>/*estβ2*<sup>1</sup> the amplified genes are 2.7 Kb apart, the difference being accounted for by expansion with three indels in the intergenic spacer (109). The indels may have introduced further regulatory elements to the amplicon intergenic spacer (52).

The identical RFLP patterns of the *estα2*<sup>1</sup> and *estβ2*<sup>1</sup> loci in resistant *C. pipiens* complex populations worldwide suggest that the amplification of these alleles occurred once and has since spread by migration (92). Other alleles of the esterase A and B loci are amplified in some *Culex* strains. For example, a strain of *C. pipiens* from Cyprus has an estimated 40 to 60 copies of the *estα5*<sup>1</sup> and *estβ5*<sup>1</sup> genes (42) whereas in the TEM-R strain from California amplification is found only at the B locus. The chromosomal region containing these esterase genes presumably represents an amplification hot spot, a theory supported by the amplification of the homologous esterase B gene in a distinct *Culex* species, *C. tritaeniorhynchus* (64).

An extensive study examining the spread and fitness of insects containing different esterase amplicons has been undertaken in southern France (93).

## Transcriptional Regulation Esterases

Elevated levels of esterases may not always be the result of gene amplification. The expression of *estα1* in the Barriol strain of *C. pipiens* from southern France is thought to be increased due to changes in an unidentified regulatory element



rather than underlying amplification of the *est* $\alpha$  gene (42). Amplified esterases can also be expressed at different levels. For example, there is fourfold more *est* $\beta$  than *est* $\alpha$  in resistant *C. quinquefasciatus*, although the genes are present in a 1:1 ratio. This difference in expression is reflected at the protein and mRNA level (63, 81a).


## Glutathione S-Transferases

The primary role of GSTs in mosquito insecticide resistance is in the metabolism of DDT to nontoxic products, although they also have a secondary role in organophosphate resistance (54). GST-based DDT resistance is common in a number of anopheline species, reflecting the heavy use of this insecticide for malaria control over several decades. Molecular characterization of GSTs is most developed in *An. gambiae*, although work on *An. dirus* from Thailand suggests that a similar arrangement of GSTs occurs. However, there is evidence for more limited allelic diversity in this species (52, 86, 87).

Two classes of insect GSTs have been recognized and members of both classes are important in the metabolism of insecticides in mosquitoes and other insects. The sequence of a single class II *An. gambiae* GST has been published (94), and we have cloned and sequenced the major class II GST from *Ae. aegypti* (D Grant, M Wajidi, H Ranson, and J Hemingway, unpublished data). This *Aedes* GST, GST-2, is over-expressed in the DDT resistant GG strain of *Ae. aegypti*. In this species the resistance mutation is thought to lead to disruption of a transacting repressor. The mutation prevents the normal function of the repressor leading to elevated levels of GST-2 enzyme in resistant mosquitoes (39).

The insect class I GSTs are encoded by a large gene family in *An. gambiae*, *M. domestica*, and *D. melanogaster*. The genomic organization of this GST class in these three insect species is strikingly different (89). In *D. melanogaster* eight divergent intronless genes are found within a 14 Kb DNA segment (106). In *An. gambiae* multiple class I GST genes are also clustered in a single location. At least one member of this family is intronless (91) but other class I GSTs in *An. gambiae* contain one or more introns (Figure 1). One of these genes, *aggst1* $\alpha$ , is alternatively spliced to produce four distinct mRNA transcripts, each of which shares a common 5' exon with a different 3' exon (89). The products of these spliced genes differ in their ability to metabolize DDT (91) and some of these metabolically active GSTs are upregulated in resistant mosquitoes (90).

The organization of this class I GST gene family in insecticide-resistant and insecticide-susceptible *An. gambiae* is very similar. Hence the actual GST-based resistance mechanism is probably caused by a *trans*-acting regulator. The development of a fine-scale microsatellite map (119) and bacterial artificial chromosome (BAC) library for *An. gambiae* have now made this species amenable to a positional cloning approach. Such an approach is being used to define the regulator responsible for this GST resistance.



**Figure 1** The class I glutathione S-transferase family of *Anopheles gambiae*, which includes an intronless gene *aggst1-2*, an alternatively spliced gene *aggst1α* with four distinct viable transcripts, and the *aggst1β* gene with two introns.

## Monooxygenases

Insect microsomal P450 monooxygenases belong to six families. Increased transcription of genes belonging to the CYP4, CYP6, and CYP9 families has been observed in insecticide-resistant strains in different insect species. As yet, it is not known which enzymes are responsible for insecticide metabolism in mosquitoes. Seventeen partial cDNAs encoding CYP4 P450s have been identified in *An. albimanus* (98) and a similar level of diversity is present in *An. gambiae* (H Ranson & F Collins, unpublished) but the role, if any, that this family plays in resistance in the mosquito is not known. Studies on the Australian cotton bollworm, *Helicoverpa armigera*, show that the resistance-associated P450s can vary between different strains; CYP6B2 is over-expressed in one pyrethroid-resistant strain (118) whereas in another, CYP4G8 is over-expressed (83). Increased expression of CYP9A1, a member of a third family, is found in the related species *Heliothis virescens* (95).

There is strong evidence to suggest that P450 monooxygenase-based resistance in *M. domestica* and *D. melanogaster* is mediated by mutations in *trans*-acting regulatory genes. CYP6A8 is highly expressed in the DDT-resistant 91-R strain of *D. melanogaster* but not detectable in the uninduced 91-C susceptible strain (28, 67). Hybrids between the two strains show low levels of expression, suggesting that the 91-C strain carries a repressor that suppresses transcription of CYP6A8. A mutation in this repressor is thought to be responsible for the high level of expression of CYP6A8 in 91-R. In the house fly, CYP6D1-mediated metabolic resistance to pyrethroids is controlled by a nearly completely dominant *cis*-factor on autosome 1 and an incompletely recessive *trans*-factor on autosome 2 (66).

## Target-Site Resistance

Non-silent point mutations within structural genes are the most common cause of target-site resistance. For selection of the mutations to occur, the resultant amino acid change must reduce the binding of the insecticide without causing a loss of primary function of the target site. Therefore the number of possible amino acid substitutions is very limited. Hence, identical resistance-associated mutations are commonly found across highly diverged taxa. The degree to which function is impaired by the resistance mutation is reflected in the fitness of resistant individuals in the absence of insecticide selection. This fitness cost has important implications for the persistence of resistance in the field.

The main sodium and GABA channel genes in insects have been cloned and their sequences compared in resistant and susceptible insects. Acetylcholinesterase-based resistance has been well characterized in *Drosophila* (36), but the elucidation of this mechanism at the molecular level in mosquitoes has proved more difficult.

## GABA Receptor Changes

The GABA receptors belong to a superfamily of neurotransmitter receptors that also includes the nicotinic acetylcholine receptors. These receptors are formed by the oligomerization of five subunits around a central transmitter-gated ion channel. Five different subunits have been cloned from vertebrates. To date only three subunits have been cloned from *Drosophila melanogaster*, but these do not fit readily into the vertebrate GABA subunit classification (61).

An alanine-to-serine substitution in the putative channel-lining domain of the GABA receptor confers resistance to cyclodienes such as dieldrin ( $\gamma$ -HCH) (34). The mutation was first identified in *Drosophila* but has since been shown to occur in a broad range of dieldrin-resistant insects, including *Ae. aegypti* (104). The only variation in resistant insects is that glycine rather than serine can sometimes be the substituted amino acid residue. Despite the widespread switch away from the use of cyclodiene insecticides for agricultural and public health use the resistance allele is still found at relatively high frequencies in insect field populations (4).

## Sodium Channels

A reduction in the sensitivity of the insect's voltage-gated sodium channels to the binding of insecticides causes the resistance phenotype known as "kdr." Changes associated with pyrethroid/DDT resistance in the sodium channels of insects are more variable than those seen in the GABA receptors but still appear to be limited to a small number of regions on this large channel protein.

The para sodium channel of houseflies contains 2108 amino acids, which fold into 4 hydrophobic repeat domains (I-IV) separated by hydrophilic linkers. The first mutation to be characterized in *kdr* insects was a leucine to phenylalanine point mutation in the S6 transmembrane segment of domain II in the sodium channel sequence of *M. domestica* (116, 117) which produces 10- to 20-fold resistance to DDT and pyrethroids. In "*super-kdr*" houseflies, this mutation also occurs with a second methionine to threonine substitution further upstream in the same domain, resulting in more than 500-fold resistance (117). Analysis of the domain region of the *para*-sodium channel gene in pyrethroid-resistant *An. gambiae* from the Ivory Coast showed an identical Leu to Phe mutation in this species (71).


A PCR-based diagnostic test discriminates between homozygous-susceptible, homozygous-resistant, and heterozygous individuals with the Leu to Phe mutation (71). Because *kdr* is semi-recessive or fully recessive (81), the ability to detect heterozygotes is of paramount importance in the early detection and management of resistance in the field.

The limited number of changes associated with *kdr*-type resistance may be constrained by the number of modifications that can influence pyrethroid/DDT binding to the sodium channels. However, a note of caution should be added:

There is already a tendency to investigate pyrethroid-resistant insects with a PCR approach confined to regions where a *kdr* mutation has already been seen. Hence consistent resistance-associated changes in other parts of the sodium-channel gene could be missed. A different approach to isolating *kdr*-type mutants has been used in *Drosophila*, utilizing the relative ease with which large numbers of mutants in the para sodium channel gene can be isolated based on their temperature sensitivity. Two classes of these mutations are in positions equivalent to the *kdr* and *super-kdr* mutations in different domains. The third class is in a novel position (Figure 2) (34).

## Acetylcholinesterase

Vertebrates have two cholinesterases: acetylcholinesterase and butyrylcholinesterase. In *D. melanogaster* only a single cholinesterase gene, *Ace*, coding for acetylcholinesterase has been cloned, based on a knowledge of its location via



**Figure 2** The location of amino acid substitutions occurring in the para-sodium channel gene of resistant strains of various insects. Mutations that have been documented in field strains of medical vectors are shown in A. B shows replacements in *Drosophila* isolated on the basis of temperature sensitivity paralytic phenotypes which confer DDT/pyrethroid resistance (adapted from 34).

isolation of a range of different mutants (43). A range of different amino acid substitutions in the *Ace* genes of *Drosophila* and the house fly *M. domestica* putatively cause resistance (reviewed in 33). Many of these resistance-associated residues are predicted to lie close to or within the acetylcholinesterase active site gorge (78). In *C. pipiens* at least two cholinesterase genes occur, both with acetylcholinesterase-like activity (69). A similar organization occurs in *Amphioxus*. To date the only acetylcholinesterase to have been cloned in *C. pipiens* is AChE2, which is not involved in insecticide resistance (69). The AChE2 gene is sex linked, unlike the resistance-conferring AChE1, which is autosomal. AChE genes have been cloned from the mosquitoes *Ae. aegypti* and *An. stephensi*, but both of these genes are also sex linked (2, 99).

As yet, there is no recorded AChE-based resistance mechanism in *An. stephensi*, but to date none of the acetylcholinesterase resistances recorded in insects have been sex linked, suggesting that these genes do not represent the insecticidal target site. However, detailed analysis of inhibition profiles for acetylcholinesterase from *Ae. aegypti* suggests that there is only one AChE locus in this species. If such is the case, altered acetylcholinesterase-based resistance in this species should be sex linked. At least one case of altered AChE has been reported in *Ae. aegypti* from Trinidad (108).

Five point mutations associated with resistance to organophosphorus and carbamate insecticides have been identified within the *D. melanogaster* acetylcholinesterase gene (78) and site-directed mutagenesis of the sex-linked AChE from *Ae. aegypti* has demonstrated that these same mutations also confer resistance in the mosquito enzyme (74). However, none of these mutations have been identified in field-collected or laboratory-selected strains of mosquitoes.

Insensitive AChE has a severe fitness cost in *C. pipiens* populations in southern France (31), which is probably caused by a reduction in the AChE activity of the mutated enzyme compared to the wild type.

## MANAGEMENT OF INSECTICIDE-RESISTANT VECTOR POPULATIONS

The practice of using an insecticide until resistance becomes a limiting factor is rapidly eroding the number of suitable insecticides for vector control. Rotations, mosaics, and mixtures have all been proposed as resistance management tools (25, 26, 96). Numerous mathematical models have been produced to estimate how these tools should be optimally used (102). However, these models have rarely been tested under field conditions for insect vectors, due to the practical difficulties in estimating changes in resistance gene frequencies in large samples of insects (56). With the advent of different biochemical and molecular techniques for resistance-gene frequency estimation, field trials of resistance management strategies have become more feasible.

A large-scale trial of the use of rotations or mosaics of insecticides compared to single use of DDT or a pyrethroid is currently underway in Mexico (56, 82). Changes in resistance gene frequencies in *An. albimanus* are being monitored over a four-year period (82). Information resulting from such large-scale trials may allow us to establish rational strategies for long-term insecticide use in disease control programs.

As our ability to manipulate the insect genome improves and our understanding of the regulation of insecticide resistance mechanisms increase, new strategies should be devised for incorporating this new knowledge into these control programs.

Visit the Annual Reviews home page at [www.AnnualReviews.org](http://www.AnnualReviews.org).

#### LITERATURE CITED

1. Amin AM, Hemingway J. 1989. Preliminary investigation of the mechanisms of DDT and pyrethroid resistance in *Culex quinquefasciatus* Say (Diptera: Culicidae) from Saudi Arabia. *Bull. Entomol. Res.* 79:361–66
2. Anthony NM, Rocheleau T, Mocelin G, Lee HJ, French-Constant RH. 1995. Cloning, sequencing and functional expression of an acetylcholinesterase gene from the yellow fever mosquito *Aedes aegypti*. *FEBS Lett.* 368:461–65
3. Ariaratnam V, Georghiou GP. 1974. Carbamate resistance in *Anopheles albimanus*. Cross-resistance spectrum and stability of resistance. *Bull. WHO* 51:655–59
4. Aronstein K, Ode P, French-Constant RH. 1995. PCR based monitoring of specific *Drosophila* (Diptera: Drosophilidae) cyclodiene resistance alleles in the presence and absence of selection. *Bull. Entomol. Res.* 85:5–9
5. Ayad H, Georghiou GP. 1975. Resistance to organophosphates and carbamates in *Anopheles albimanus* based on reduced sensitivity to acetylcholinesterase. *J. Econ. Entomol.* 68:295–97
6. Bang YH, Tonn RJ, Punurai P. 1969. Insecticide susceptibility and resistance found in 14 strains of *Aedes aegypti* collected from Bangkok-Thonburi, Thailand. *WHO/Vector Biol. Control* 69:117
7. Bencheikh H, BenAliHaouas Z, Marquine M, Pasteur N. 1998. Resistance to organophosphorus and pyrethroid insecticides in *Culex pipiens* (Diptera: Culicidae) from Tunisia. *J. Med. Entomol.* 35:251–60
8. Berge JB, Feyereisen R, Amichot M. 1998. Cytochrome P450 monooxygenases and insecticide resistance in insects. *Philos. Trans. R. Soc. London Ser. B* 353:1701–5
9. Bermudez I, Hawkins CA, Taylor AM, Beadle DJ. 1991. Actions of insecticides on the insect GABA receptor complex. *J. Recept. Res.* 11:221–32
10. Bloomquist JR. 1994. Cyclodiene resistance at the insect GABA receptor chloride channel complex confers broad cross-resistance to convulsants and experimental phenylpyrazole insecticides. *Arch. Insect Biochem. Physiol.* 26:69–79
11. Boddington RG. 1992. Characterization of malathion carboxylesterases and non-specific esterases in the malaria mosquito *Anopheles stephensi*. PhD diss. Univ. London
12. Bradley DJ. 1998. The particular and general. Issues of specificity and verti-

- cality in the history of malaria control. *Parassitologia* 40:5–10
13. Brogdon WG, Barber AM. 1990. Fenitrothion-deltamethrin cross-resistance conferred by esterases in Guatemalan *Anopheles albimanus*. *Pest. Biochem. Physiol.* 37:130–39
  14. Brogdon WG, McAllister JC, Vulule J. 1997. Heme peroxidase activity measured in single mosquitoes identifies individuals expressing the elevated oxidase mechanism for insecticide resistance. *J. Am. Mosq. Control Assoc.* 13:233–37
  15. Brown AWA. 1986. Insecticide resistance in mosquitoes: a pragmatic review. *J. Am. Mosq. Control Assoc.* 2:123–40
  16. Brown AWA, Pal R. 1971. Insecticide resistance in arthropods. *WHO Monogr. Ser.* 38
  17. Campbell PM, Newcomb RD, Russell RJ, Oakeshott JG. 1998. Two different amino acid substitutions in the al-esterase, E3, confer alternative types of organophosphorus insecticide resistance in the sheep blowfly, *Lucilia cuprina*. *Insect Biochem. Mol. Biol.* 28:139–50
  18. Chandre F, Darriet F, Brengues C, Manguin S, Carnevale P, Guillet P. 1999. Pyrethroid cross-resistance spectrum among populations of *Anopheles gambiae* from Cote D'Ivoire. *J. Am. Mosq. Control Assoc.* 15:53–59
  19. Chandre F, Darriet F, Darder M, Cuany A, Doannio JMC, et al. 1998. Pyrethroid resistance in *Culex quinquefasciatus* from West Africa. *Med. Vet. Entomol.* 12:359–66
  20. Chandre F, Darriet F, Manga L, Akogbeto M, Faye O, et al. 1999. Status of pyrethroid resistance in *Anopheles gambiae* s.l. *Bull. WHO* 77:230–34
  21. Chosidow O, Chastang C, Brue C, Bouvet E, Izri M, et al. 1995. Controlled study of malathion and *d*-phenothrin lotions for *Pediculus humanus* var *capitis*-infested schoolchildren. *Lancet* 345:731–32
  22. Clark AG, Dick GL, Martindale SM, Smith JN. 1985. Glutathione S-transferases from the New Zealand grass grub, *Costelytra zealandica*. *Insect Biochem.* 15:35–44
  23. Clark AG, Shamaan NA. 1984. Evidence that DDT-dehydrochlorinase from the house fly is a glutathione S-transferase. *Pestic. Biochem. Physiol.* 22:249–61
  24. Clark AG, Shamaan NA, Dauterman WC, Hayaoka T. 1984. Characterization of multiple glutathione transferases from the housefly, *Musca domestica* (L). *Pestic. Biochem. Physiol.* 22:51–59
  25. Curtis CF. 1985. Theoretical models of the use of insecticide mixtures for the management of resistance. *Bull. Entomol. Res.* 75:259–65
  26. Curtis CF, Hill N, Kasim SH. 1993. Are there effective resistance management strategies for vectors of human disease? *Biol. J. Linn. Soc.* 48:3–18
  27. DeSilva D, Hemingway J, Ranson H, Vaughan A. 1997. Resistance to insecticides in insect vectors of disease: Est $\alpha$ 3, a novel amplified esterase associated with est $\alpha$ 1s from insecticide resistant strains of the mosquito *Culex quinquefasciatus*. *Exp. Parasitol.* 87:253–59
  28. Dombrowski SM, Krishnan R, Witte M, Maitra S, Diesing C, et al. 1998. Constitutive and barbital-induced expression of the CYP6A2 allele of a high producer strain of CYP6A2 in the genetic background of a low producer strain. *Gene* 221:69–77
  29. El-Sayed S, Hemingway J, Lane RP. 1989. Susceptibility baselines for DDT metabolism and related enzyme systems in the sandfly *Phlebotomus papatasi* (Scopoli) (Diptera:Psychodidae). *Bull. Entomol. Res.* 79:679–84
  30. Eshgy N. 1978. Tolerance of *Anopheles stephensi* to malathion in the province of Fars, Southern Iran 1977. *Mosq. News* 38:580–83
  31. Evgenev MB, Corces VG, Lankenau DH. 1992. Ulysses transposable element



- of *Drosophila* shows high structural similarities to functional domains of retroviruses. *J. Mol. Biol.* 225:917–24
32. Farnham AW, Sawicki RM. 1976. Development of resistance to pyrethroids in insects resistant to other insecticides. *Pestic. Sci.* 7:278–82
  33. Feyereisen R. 1995. Molecular biology of insecticide resistance. *Toxicol. Lett.* 82:83–90
  34. French-Constant RH, Pittendrigh B, Vaughan A, Anthony N. 1998. Why are there so few resistance-associated mutations in insecticide target genes? *Philos. Trans. R. Soc. London Ser. B* 353:1685–93
  35. Fine BC. 1963. The present status of resistance to pyrethroid insecticides. *Pyrethrum Post* 7:18–21
  36. Fournier D, Mutero A, Pralavorio M, Bride J-M. 1993. *Drosophila* acetylcholinesterase: mechanisms of resistance to organophosphates. *Chem.-Biol. Interact.* 87:233–38
  37. Grant DF. 1991. Evolution of glutathione S-transferase subunits in Culicidae and related Nematocera: Electrophoretic and immunological evidence for conserved enzyme structure and expression. *Insect Biochem.* 21:435–45
  38. Grant DF, Dietze EC, Hammock BD. 1991. Glutathione S-transferase isozymes in *Aedes aegypti*: purification, characterization, and isozyme specific regulation. *Insect. Biochem.* 4:421–33
  39. Grant DF, Hammock BD. 1992. Genetic and molecular evidence for a trans-acting regulatory locus controlling glutathione S-transferase-2 expression in *Aedes aegypti*. *Mol. Gen. Genet.* 234:169–76
  40. Grant DF, Matsumura F. 1988. Glutathione S-transferase-1 in *Aedes aegypti* larvae. Purification and properties. *Insect Biochem.* 18:615–22
  41. Grant DF, Matsumura F. 1989. Glutathione S-transferase 1 and 2 in susceptible and insecticide resistant *Aedes aegypti*. *Pestic. Biochem. Physiol.* 33:132–43
  42. Guillemaud T, Makate N, Raymond M, Hirst B, Callaghan A. 1997. Esterase gene amplification in *Culex pipiens*. *Insect Mol. Biol.* 6:319–27
  43. Hall LMC, Spierer P. 1986. The Ace locus of *Drosophila melanogaster*: structural gene for acetylcholinesterase with an unusual 5' leader. *EMBO J.* 5:2949–54
  44. Hemingway J. 1982. The biochemical nature of malathion resistance in *Anopheles stephensi* from Pakistan. *Pestic. Biochem. Physiol.* 17:149–55
  45. Hemingway J. 1983. Biochemical studies on malathion resistance in *Anopheles arabiensis* from Sudan. *Trans. R. Soc. Trop. Med. Hyg.* 77:477–80
  46. Hemingway J. 1985. Malathion carboxylesterase enzymes in *Anopheles arabiensis* from Sudan. *Pest. Biochem. Physiol.* 23:309–13
  47. Hemingway J. 1999. Insecticide resistance in malaria vectors: A new approach to an old subject. *Parassitologia*. In press
  48. Hemingway J, Boddington RG, Harris J, Dunbar SJ. 1989. Mechanisms of insecticide resistance in *Aedes aegypti* (L.) (Diptera:Culicidae) from Puerto Rico. *Bull. Entomol. Res.* 79:123–30
  49. Hemingway J, Callaghan A, Kurtak DC. 1991. Biochemical characterization of chlorphoxim resistance in adults and larvae of the *Simulium damnosum* complex (Diptera: Simuliidae). *Bull. Entomol. Res.* 81:401–6
  50. Hemingway J, Coleman M, Vaughan A, Patton M, DeSilva D. 1999. Aldehyde oxidase is co-amplified with the worlds most common *Culex* mosquito insecticide resistance-associated esterases. *Insect Mol. Biol.* In press
  51. Hemingway J, Georghiou GP. 1983. Studies on the acetylcholinesterase of *Anopheles albimanus* resistant and susceptible to organophosphate and carbamate insecticides. *Pestic. Biochem. Physiol.* 19:167–71
  52. Hemingway J, Hawkes N, Prapanthadara

- L, Jayawardena KGI, Ranson H. 1998. The role of gene splicing, gene amplification and regulation in mosquito insecticide resistance. *Philos. Trans. R. Soc. London Ser. B* 353:1695–99
53. Hemingway J, Karunaratne SHPP. 1998. Mosquito carboxylesterases: a review of the molecular biology and biochemistry of a major insecticide resistance mechanism. *Med. Vet. Entomol.* 12:1–12
54. Hemingway J, Malcolm CA, Kissoon KE, Boddington RG, Curtis CF, Hill N. 1985. The biochemistry of insecticide resistance in *Anopheles sacharovi*: comparative studies with a range of insecticide susceptible and resistant *Anopheles* and *Culex* species. *Pestic. Biochem. Physiol.* 24:68–76
55. Hemingway J, Miyamoto J, Herath PRJ. 1991. A possible novel link between organo phosphorus and DDT insecticide resistance genes in *Anopheles*: supporting evidence from fenitrothion metabolism studies. *Pestic. Biochem. Physiol.* 39:49–56
56. Hemingway J, Penilla RP, Rodriguez AD, James BM, Edge W, et al. 1997. Resistance management strategies in malaria vector mosquito control. A large-scale trial in Southern Mexico. *Pestic. Sci.* 51:375–82
57. Hemingway J, Small GJ, Monro A, Sawyer BV, Kasap H. 1992. Insecticide resistance gene frequencies in *Anopheles sacharovi* populations of the Cukurova plain, Adana province, Turkey. *Med. Vet. Entomol.* 6:342–48
58. Herath PRJ, Davidson G. 1981. Multiple resistance in *Anopheles culicifacies* Giles. *Mosq. News* 41:325–27
59. Herath PRJ, Hemingway J, Weerasinghe IS, Jayawardena KGI. 1987. The detection and characterization of malathion resistance in field populations of *Anopheles culicifacies* B in Sri Lanka. *Pestic. Biochem. Physiol.* 29:157–62
60. Herath PRJ, Miles SJ, Davidson G. 1981. Fenitrothion (OMS 43) resistance in the taxon *Anopheles culicifacies* Giles. *J. Trop. Med. Hyg.* 84:87–88
61. Hosie AM, Aronstein K, Sattelle DB, French-Constant RH. 1997. Molecular biology of insect neuronal GABA receptors. *Trends Neurosci.* 20:578–83
62. Kadous AA, Ghiasuddin SM, Matsu-mura F, Scott JG, Tanaka K. 1983. Difference in the picrotoxinin receptor between the cyclodiene-resistant and susceptible strains of the German-cockroach. *Pest. Biochem. Physiol.* 19:157–66
63. Karunaratne SHPP, Hemingway J, Jayawardena KGI, Dassanayaka V, Vaughan A. 1995. Kinetic and molecular differences in the amplified and non-amplified esterases from insecticide-resistant and susceptible *Culex quinquefasciatus* mosquitoes. *J. Biol. Chem.* 270:31124–28
64. Karunaratne SHPP, Vaughan A, Paton MG, Hemingway J. 1998. Amplification of a serine esterase gene is involved in insecticide resistance in Sri Lankan *Culex tritaeniorhynchus*. *Insect Mol. Biol.* 7:307–15
65. Kasai S, Weerasinghe IS, Shono T. 1998. P<sup>450</sup> Monooxygenases are an important mechanism of permethrin resistance in *Culex quinquefasciatus* say larvae. *Arch. Insect Biochem. Physiol.* 37:47–56
66. Liu N, Scott JG. 1997. Inheritance of CYP6D1-mediated pyrethroid resistance in house fly (Diptera: Muscidae). *J. Econ. Entomol.* 90:1478–81
67. Maitra S, Dombrowski SM, Waters LC, Gunguly R. 1996. Three second chromosome-linked clustered CYP6 genes show differential constitutive and barbital-induced expression in DDT-resistant and susceptible strains of *Drosophila melanogaster*. *Gene* 180:165–71
68. Malcolm CA, Boddington RG. 1989. Malathion resistance conferred by a carboxylesterase in *Anopheles culicifacies* Giles (Species B) (Diptera: Culicidae). *Bull. Entomol. Res.* 79:193–99
69. Malcolm CA, Bourguet D, Ascolillo A,

- Rooker SJ, Garvey CF, et al. 1998. A sex-linked *Ace* gene, not linked to insensitive acetylcholinesterase-mediated insecticide resistance in *Culex pipiens*. *Insect Mol. Biol.* 7:107–20
70. Malcolm CA, Wood RJ. 1982. Location of a gene conferring resistance to knock-down by permethrin and bioresmethrin in adults of the BKPM3 strain of *Aedes aegypti*. *Genetica* 59:233–37
71. Martinez-Torres D, Chandre F, Williamson MS, Darriet F, Berge JB, et al. 1998. Molecular characterization of pyrethroid knockdown resistance (*kdr*) in the major malaria vector *Anopheles gambiae* s.s. *Insect Mol. Biol.* 7:179–84
72. Matsumura F, Brown AWA. 1961. Biochemical study of malathion tolerant *Aedes aegypti*. *Mosq. News* 21:192–94
73. Maunder JW. 1971. Resistance to organochlorine insecticides in head lice and trials using alternative compounds. *Med. Off.* 258:27–31
74. Milatovic D, Moretto A, Osman KA, Lotti M. 1997. Phenyl valerate esterases other than neuropathy target esterase and the promotion of organophosphate polyneuropathy. *Chem. Res. Toxicol.* 10:1045–48
75. Mouches C, Pasteur N, Berge JB, Hyrien O, Raymond M, et al. 1986. Amplification of an esterase gene is responsible for insecticide resistance in a Californian *Culex* mosquito. *Science* 233:778–80
76. Mourya DT, Hemingway J, Leake CJ. 1993. Changes in enzyme titres with age in four geographical strains of *Aedes aegypti* and their association with insecticide resistance. *Med. Vet. Entomol.* 7:11–16
77. Mumcuoglu KY, Miller J, Uspensky I, Hemingway J, Klaus S, et al. 1995. Pyrethroid resistance in the head louse *Pediculus humanus capitis* from Israel. *Med. Vet. Entomol.* 9:427–32
78. Mutero A, Pralavorio M, Bride JM, Fournier D. 1994. Resistance-associated point mutations in insecticide-insensitive acetylcholinesterase. *Proc. Natl. Acad. Sci. USA* 91:5922–26
79. Newcomb RD, Campbell PM, Russell RJ, Oakeshott JG. 1997. cDNA cloning, baculovirus-expression and kinetic properties of the esterase, E3, involved in organophosphorus resistance in *Lucilia cuprina*. *Insect Biochem. Mol. Biol.* 27:15–25
80. Newcomb RD, East PD, Russell RJ, Oakeshott JG. 1996. Isolation of  $\alpha$  cluster esterase genes associated with organophosphate resistance in *Lucilia cuprina*. *Insect Mol. Biol.* 5:211–16
81. Ozers MS, Friesen PD. 1996. The *Env*-like open reading frame of the baculovirus-integrated retrotransposon *ted* encodes a retrovirus-like envelope protein. *Virology* 226:252–59
- 81a. Paton MG, Karunaratne SHPP, Giakoumaki E, Roberts N, Hemingway J. 2000. Quantitative analysis of gene amplification in insecticide resistant *Culex* mosquitoes. *Biochem. J.* In press
82. Penilla RP, Rodrigues AD, Hemingway J, Torres JL, Arredondo-Jimenez JJ, Rodriguez MH. 1998. Resistance management strategies in malaria vector mosquito control. Baseline data for a large-scale field trial against *Anopheles albimanus* in Mexico. *Med. Vet. Entomol.* 12:217–33
83. Pittendrigh B, Aronstein K, Zinkovskiy E, Andreev O, Campbell BC, et al. 1997. Cytochrome P450 genes from *Helicoverpa armigera*: expression in a pyrethroid-susceptible and -resistant strain. *Insect Biochem. Mol. Biol.* 27:507–12
84. Prapanthadara L, Hemingway J, Ketterman AJ. 1993. Partial purification and characterization of glutathione S-transferase involved in DDT resistance from the mosquito *Anopheles gambiae*. *Pest. Biochem. Physiol.* 47:119–33
85. Prapanthadara L, Hemingway J, Ketterman AJ. 1995. DDT-resistance in *Anopheles gambiae* Giles from Zanzibar Tanzania, based on increased DDT-dehy-

- drochlorinase activity of glutathione S-transferases. *Bull. Entomol. Res.* 85:267–74
86. Prapanthadara L, Koottathep S, Promtet N, Hemingway J, Ketterman AJ. 1996. Purification and characterization of a major glutathione S-transferase from the mosquito *Anopheles dirus* (Species B). *Insect Biochem. Mol. Biol.* 26:277–85
87. Prapanthadara LA, Ranson H, Somboon P, Hemingway J. 1998. Cloning, expression and characterization of an insect class I glutathione S-transferase from *Anopheles dirus* species B. *Insect Biochem. Mol. Biol.* 28:321–29
88. Raghavendra K, Vasantha K, Subbarao SK, Pillai MKK, Sharma VP. 1991. Resistance in *Anopheles culicifacies* sibling species B and C to malathion in Andhra Pradesh and Gujarat states, India. *J. Am. Mosq. Control Assoc.* 7:255–59
89. Ranson H, Collins FH, Hemingway J. 1998. The role of alternative mRNA splicing in generating heterogeneity within the *Anopheles gambiae* class I glutathione S-transferase family. *Proc. Natl. Acad. Sci. USA* 95:14284–89
90. Ranson H, Cornel AJ, Fournier D, Vaughan A, Collins FH, Hemingway J. 1997. Cloning and localization of a glutathione S-transferase class I gene from *Anopheles gambiae*. *J. Biol. Chem.* 272:5464–68
91. Ranson H, Prapanthadara L, Hemingway J. 1997. Cloning and characterisation of two glutathione S-transferases from a DDT resistant strain of *Anopheles gambiae*. *Biochem. J.* 324:97–102
92. Raymond M, Callaghan A, Fort P, Pasteur N. 1991. Worldwide migration of amplified insecticide resistance genes in mosquitoes. *Nature* 350:151–53
93. Raymond M, Chevillon C, Guillemaud T, Lenormand T, Pasteur N. 1998. An overview of the evolution of overproduced esterases in the mosquito *Culex pipiens*. *Philos. Trans. R. Soc. London Ser. B* 353:1707–11
94. Reiss RA, James AA. 1993. A glutathione S-transferase gene of the vector mosquito, *Anopheles gambiae*. *Insect Mol. Biol.* 2:25–32
95. Rose RL, Goh D, Thompson DM, Verma KD, Heckel DG, et al. 1997. Cytochrome P450 (CYP)9A1 in *Heliothis virescens*: the first member of a new CYP family. *Insect Biochem. Mol. Biol.* 27:605–15
96. Roush RT. 1989. Designing resistance management programmes: how can you choose? *Pestic. Sci.* 26:423–42
97. Rupes V, Moravec J, Chmela J, Ledvidka J, Zelenkova J. 1994. A resistance of head lice (*Pediculus capitis*) to permethrin in Czech republic. *Centr. Eur. J. Public Health* 3:30–32
98. Scott JA, Collins FH, Feyereisen R. 1994. Diversity of cytochrome P<sup>450</sup> genes in the mosquito, *Anopheles albimanus*. *Biochem. Biophys. Res. Commun.* 205:1452–59
99. Severson DW, Anthony NM, Andreen O, French-Constant RH. 1997. Molecular mapping of insecticide resistance genes in the yellow fever mosquito (*Aedes aegypti*). *J. Hered.* 88:520–24
100. Soderlund DM, Bloomquist JR. 1989. Neurotoxic action of pyrethroid insecticides. *Annu. Rev. Entomol.* 34:77–96
101. Subbarao SK, Vasantha K, Joshi H, Raghavendra K, Devi CU, et al. 1992. Role of *Anopheles culicifacies* sibling species in malaria transmission in Madhya Pradesh state, India. *Trans. R. Soc. Trop. Med. Hyg.* 86:613–14
102. Tabashnik BE. 1989. Managing resistance with multiple pesticide tactics: theory, evidence and recommendations. *J. Econ. Entomol.* 82:1263–69
103. Taplin D, Meinking TL. 1987. Pyrethrins and pyrethroids for the treatment of scabies and pediculosis. *Semin. Dermatol.* 6:125–35
104. Thompson M, Shotkoski F, French-Constant RH. 1993. Cloning and sequencing of the cyclodiene insecticide resistance gene from the yellow fever

- mosquito *Aedes aegypti*. *FEBS Lett.* 325:187–90
105. Toung YS, Hsieh T, Tu CD. 1990. *Drosophila* glutathione S-transferase 1-1 shares a region of sequence homology with maize glutathione S-transferase III. *Proc. Natl. Acad. Sci. USA* 87:31–35
106. Toung YS, Hsieh T, Tu CD. 1993. The glutathione S-transferase *D* genes: a divergently organized, intronless gene family in *Drosophila melanogaster*. *J. Biol. Chem.* 268:9737–46
107. Vatandoost H, McCaffery AR, Townson H. 1996. An electrophysiological investigation of target site insensitivity mechanisms in permethrin-resistant and susceptible strains of *Anopheles stephensi*. *Trans. R. Soc. Trop. Med. Hyg.* 90:216
108. Vaughan A, Chadee DD, French-Constant RH. 1998. Biochemical monitoring of organophosphorus and carbamate insecticide resistance in *Aedes aegypti* mosquitoes from Trinidad. *Med. Vet. Entomol.* 12:318–21
109. Vaughan A, Hawkes N, Hemingway J. 1997. Co-amplification explains linkage disequilibrium of two mosquito esterase genes in insecticide-resistant *Culex quinquefasciatus*. *Biochem. J.* 325:359–65
110. Vaughan A, Hemingway J. 1995. Mosquito carboxylesterase Est $\alpha$ 2<sup>1</sup> (A2). Cloning and sequence of the full length cDNA for a major insecticide resistance gene worldwide in the mosquito *Culex quinquefasciatus*. *J. Biol. Chem.* 270:17044–49
111. Vaughan A, Rodriguez M, Hemingway J. 1995. The independent gene amplification of indistinguishable esterase B electromorphs from the insecticide resistant mosquito *Culex quinquefasciatus*. *Biochem. J.* 305:651–58
112. Vulule JM, Beach RF, Atieli FK, Roberts JM, Mount DL, Mwangi RW. 1994. Reduced susceptibility of *Anopheles gambiae* to permethrin associated with the use of permethrin-impregnated bed-nets and curtains in Kenya. *Med. Vet. Entomol.* 8:71–75
113. WHO. 1992. Vector resistance to pesticides. Fifteenth report of the expert committee on vector biology and control. In *WHO Tech. Rep. Ser.* 818:1–55
114. Whyard S, Downe AER, Walker VK. 1994. Isolation of an esterase conferring insecticide resistance in the mosquito *Culex tarsalis*. *Insect Biochem. Mol. Biol.* 24:819–27
115. Whyard S, Downe AER, Walker VK. 1995. Characterization of a novel esterase conferring insecticide resistance in the mosquito *Culex tarsalis*. *Arch. Insect Biochem. Physiol.* 29:329–42
116. Williamson MS, Denholm I, Bell CA, Devonshire AL. 1993. Knockdown resistance (kdr) to DDT and pyrethroid insecticides maps to a sodium channel gene locus in the housefly (*Musca domestica*). *Mol. Gen. Genet.* 240:17–22
117. Williamson MS, Martinez-Torres D, Hick CA, Devonshire AL. 1996. Identification of mutations in the housefly para-type sodium channel gene associated with knockdown resistance (kdr) to pyrethroid insecticides. *Mol. Gen. Genet.* 252:51–60
118. Xiao-Ping W, Hobbs AA. 1995. Isolation and sequence analysis of a cDNA clone for a pyrethroid inducible P450 from *Helicoverpa amrigera*. *Insect Biochem. Mol. Biol.* 25:1001–9
119. Zheng L, Benedict MQ, Cornel AJ, Collins FH, Kafatos FC. 1996. An integrated genetic map of the African human malaria vector mosquito *Anopheles gambiae*. *Genetics* 143:941–45
120. Ziegler R, Whyard S, Downe AER, Wyatt GR, Walker VK. 1987. General esterase, malathion carboxylesterase, and malathion resistance in *Culex tarsalis*. *Pestic. Biochem. Physiol.* 8:279–85

